

The Darboux Transform and some Integrable cases of the q -Riccati Equation¹

Anatol Odziejewicz* & Alina Ryzko**

Institute of Theoretical Physics
University in Białystok
ul. Lipowa 41, 15-424 Białystok, Poland
E-mail: *aodziejew@labfiz.uwb.edu.pl, **alaryzko@alpha.uwb.edu.pl

Abstract

Using the q -version of the Darboux transform we obtain the general solution of q -difference Riccati equation from a special one by the action of one-parameter group. This allows us to construct the solutions for the large class of q -difference Riccati equations as well as q -difference Schrödinger equations.

Introduction

In this paper we investigate the Darboux factorization method for q -difference version of Riccati and Schrödinger equations. It appears that this method which is by all means effective for differential Riccati and Schrödinger equation ([7, 8, 2]) leads to non-trivial and interesting results in the q -deformed case too. Some of the new formulae have their undeformed version. They tend in the limit of $q \rightarrow 1$ to the ones, which are well known in differential case.

The Darboux factorization

$$-\frac{d^2}{dx^2} + V(x) = \left(\frac{d}{dx} + u(x) \right) \left(-\frac{d}{dx} + u(x) \right) \quad (1)$$

gives the well known correspondence between one-dimensional Schrödinger equation

$$\left(-\frac{d^2}{dx^2} \psi(x) + V(x) \right) \psi(x) = 0 \quad (2)$$

and the Riccati equation

$$\frac{d}{dx} u(x) = -u^2(x) + V(x), \quad (3)$$

¹ Supported in part by KBN grant 2 PO3 A 012 19

where

$$u(x) = \frac{d}{dx} \ln \psi(x). \quad (4)$$

This correspondence is a starting point for a search of the exact solutions of both equations above (see [2]).

We will investigate, in this paper, an analogue of the Darboux method for the pair of q -difference equation

$$\partial_q \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \begin{pmatrix} R(x) & S(x) \\ V(x) & T(x) \end{pmatrix} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix}, \quad (5)$$

and

$$V(x) = \partial_q u(x) - T(x)u(x) + R(x)u(qx) + S(x)u(x)u(qx). \quad (6)$$

whose solutions are related by

$$u(x) = \frac{\varphi(x)}{\psi(x)}. \quad (7)$$

It is clear that (5) and (6) generalize (2) and (3) respectively. The Schrödinger and Riccati equations are obtained in the limit $q \rightarrow 1$ under additional assumption that $R(x) = 0 = T(x)$.

Let us recall here, that the q -derivative and q -integral are defined by

$$\partial_q \psi(x) = \frac{\psi(x) - \psi(qx)}{(1-q)x}, \quad (8)$$

$$\int_0^x \psi(x) d_q t = \sum_{n=0}^{\infty} (1-q) q^n x \psi(q^n x) \quad (9)$$

respectively, where $0 \leq q \leq 1$. The standard derivative and integral are obtained for $q = 1$. However, the reason for the investigation of the q -difference equations (5) and (6) is not only that they generalize in a natural way the Schrödinger and Riccati equations.

If one additionally assumes $1 - (1-q)xT(x) = 0$ in the real case and one takes q^n instead of the real argument x , the equation (5) appear to reduce to the three terms recurrence equation

$$\psi_{n+2} = [1 - (1-q)q^{n+1}R(q^{n+1})] \psi_{n+1} + (1-q)^2 q^{2n+1} S(q^{n+1}) V(q^n) \psi_n, \quad (10)$$

for the function $\psi_n := \psi(q^n)$ of the natural argument $n \in \mathbb{N} \cup \{0\}$. Hence, the q -difference equation (5) can be applied to those physical problems which are related to the theory of orthogonal polynomials (see [4]).

The paper is organized in the following way. In Section 1 we will introduce the q -difference Darboux transform and we will integrate the equation (5) for the case of $V(x) = 0$. The action of q -difference Darboux transform will be presented in Section 2. There we find an one-parameter auto-Bäcklund transform for the q -difference Riccati equation and we show that it generates the general solution of (6) from a special one. Section 3 is devoted to the presentation of some extended classes of solutions of the q -difference Schrödinger and Riccati equation. All results presented in Section 1 and 2 have well known differential counterparts and this aspect is also exhibited in the paper.

1 The q -difference Darboux transform

In order to solve the q -difference equation (5) by the iterative method we will rewrite it in the following form

$$\begin{pmatrix} \psi(qx) \\ \varphi(qx) \end{pmatrix} = \Lambda(x) \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix}, \quad (11)$$

where

$$\Lambda(x) = \mathbf{1} - (1 - q)x \begin{pmatrix} R(x) & S(x) \\ V(x) & T(x) \end{pmatrix}. \quad (12)$$

Let us assume here that $R(x)$, $S(x)$, $V(x)$ and $T(x)$ are continuous functions of a real argument. Hence, the matrix sequence

$$\Lambda(q^{n-1}x) \dots \Lambda(qx) \Lambda(x) =: \Lambda(x; q)_n \quad (13)$$

is pointwise convergent

$$\Lambda(x; q)_n \longrightarrow_{n \rightarrow \infty} \Lambda(x; q)_\infty \quad (14)$$

to a matrix function $\Lambda(x; q)_\infty$. The inverse matrix function $\Lambda(x; q)_\infty^{-1}$ is exactly the resolvent of equation (5), i. e.

$$\begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \Lambda(x; q)_\infty^{-1} \begin{pmatrix} \psi(0) \\ \varphi(0) \end{pmatrix}. \quad (15)$$

The problem of solving (5) thus is equivalent to the calculation of the infinite matrix product

$$\Lambda(x; q)_\infty := \prod_{k=0}^{\infty} \Lambda(q^k x). \quad (16)$$

The above suggests the following transform:

$$\Lambda(x) \longrightarrow D(qx)^{-1} \Lambda(x) D(x) = \Lambda'(x), \quad (17)$$

$$\begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} \longrightarrow D(x)^{-1} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \begin{pmatrix} \psi'(x) \\ \varphi'(x) \end{pmatrix}, \quad (18)$$

where $D(x)$ is $GL(2, \mathbb{R})$ -valued function of the real argument. It is obvious that the transform (17-18) preserves the form of the equation (11) and the transformed resolvent $\Lambda'(x; q)_\infty^{-1}$ is related to the initial one by

$$\Lambda'(x; q)_\infty^{-1} = D(x)^{-1} \Lambda(x; q)_\infty^{-1} D(0). \quad (19)$$

Thus the virtue of the above transform is to find such matrix valued function $D(x)$ for equation (5), which reduces unknown resolvent $\Lambda(x; q)_\infty^{-1}$ to some known one $\Lambda'(x; q)_\infty^{-1}$.

We will find later the explicit form of the resolvent $\Lambda(x; q)_\infty$ in the case when $V(x) = 0$. So, in order to integrate (5) it is enough to transform (16) to the uppertriangular matrix function $\Lambda(x)$ by the use of (17-18). Any matrix can be decomposed generically into the product of the uppertriangular and lowertriangular matrices. Thus, without loss of generality we can assume that

$$D(x) = \begin{pmatrix} 1 & 0 \\ -c(x) & 1 \end{pmatrix}. \quad (20)$$

After substituting (20) into (17) we find that $\Lambda'(x)$ will be an uppertriangular matrix if and only if the function $c(x)$ satisfies the q -difference Riccati equation (6).

The q -difference Schrödinger operator factorizes into the form:

$$-\partial_q^2 + V(x) = (\partial_q + u(qx)) (-\partial_q + u(x)), \quad (21)$$

iff the function $u(x)$ satisfies the equation (6) with $R(x) = T(x) = 0$ and $S(x) = 1$. Hence, it is natural to call the matrix transform (17-18) **the q -difference Darboux transform**.

We will use the identity

$$\prod_{n=0}^{\infty} (1 - (1-q)q^n x f(q^n x)) = \exp \left(\frac{1}{1-q} \int_0^x \frac{\ln(1 - (1-q)t f(t))}{t} d_q t \right), \quad (22)$$

which is an easy consequence of the definition of q -integral.

Proposition 1 *If $R(x)$, $S(x)$ and $T(x)$ are continuous functions and $V(x) = 0$ then*

$$\Lambda(x; q)_{\infty} = \begin{bmatrix} \exp \frac{1}{1-q} \int_0^x \frac{\ln(1-(1-q)tR(t))}{t} d_q t & B(x) \\ 0 & \exp \frac{1}{1-q} \int_0^x \frac{\ln(1-(1-q)tT(t))}{t} d_q t \end{bmatrix}, \quad (23)$$

where

$$B(x) = \left[-\exp \left(\frac{1}{1-q} \int_0^x \frac{\ln(1 - (1-q)tT(t))}{t} d_q t \right) \right] \times \\ \times \left[\int_0^x \frac{S(t)}{1 - (1-q)tR(t)} \exp \left(\frac{1}{1-q} \int_0^t \frac{1}{s} \ln \frac{1 - (1-q)sR(s)}{1 - (1-q)sT(s)} d_q s \right) d_q t \right]. \quad (24)$$

Proof:

Since $V(x) = 0$ we can assume that $\Lambda(x; q)_{\infty}$ is the uppertriangular matrix of the form

$$\Lambda(x; q)_{\infty} = \begin{bmatrix} \prod_{n=0}^{\infty} (1 - (1-q)q^n x R(q^n x)) & B(x) \\ 0 & \prod_{n=0}^{\infty} (1 - (1-q)q^n x T(q^n x)) \end{bmatrix}. \quad (25)$$

where from the equation

$$\Lambda(x; q)_{\infty} = \Lambda(qx; q)_{\infty} \Lambda(x), \quad (26)$$

we find that the function $B(x)$ does satisfy

$$B(x) = [1 - (1-q)xT(x)] B(qx) - \prod_{n=0}^{\infty} (1 - (1-q)q^{n+1}xR(q^{n+1}x)) (1-q)xS(x). \quad (27)$$

The equation (27) is solved by iterative method. One gets finally:

$$B(x) = - \prod_{n=0}^{\infty} (1 - (1-q)q^n x T(q^n x)) \times \\ \times \sum_{n=0}^{\infty} \frac{(1-q)q^n x S(q^n x)}{1 - (1-q)q^n x R(q^n x)} \prod_{k=n}^{\infty} \frac{(1 - (1-q)q^k x R(q^k x))}{(1 - (1-q)q^k x T(q^k x))}. \quad (28)$$

Substituting (28) into (25) and using identity (22) we obtain the formulae (23) and (24).

QED

2 The solution of q -difference Schrödinger equation and auto-Bäcklund transform for q -difference Riccati equation

We have obtained in Section 1, the resolvent function $\Lambda(x; q)_{\infty}^{-1}$ for the case of $V(x) = 0$, (see **Proposition 1**). Let us assume $\Lambda'(x)$ in (17) to be uppertriangular matrix function. Applying the q -difference Darboux transform to $\Lambda(x)$, with $D(x)$ given by (20), we find the general solution of the q -differential equation (5).

Proposition 2 *Let us assume that functions $R(x)$, $S(x)$, $T(x)$ and $V(x)$ from the equations (5) are continuous. Let $(\psi_0(x) \ \varphi_0(x))^\top$ be a particular solution of (5) one and put*

$$u_0(x) = \frac{\varphi_0(x)}{\psi_0(x)}. \quad (29)$$

Then the potential $V(x)$ is given by

$$V(x) = \partial_q u_0(x) - T(x)u_0(x) + R(x)u_0(qx) + S(x)u_0(x)u_0(qx), \quad (30)$$

and the general solutions of (5) is given by:

$$\begin{aligned} \psi(x) = & \exp \left(-\frac{1}{1-q} \int_0^x \frac{\ln \{1 - (1-q)t [R(t) + u_0(t)S(t)]\}}{t} d_q t \right) \times \\ & \times \left[D + F \int_0^x \frac{S(t)}{1 - (1-q)t [R(t) + u_0(t)S(t)]} \times \right. \\ & \times \exp \left(\frac{1}{1-q} \int_0^t \frac{1}{s} \ln \frac{1 - (1-q)s [R(s) + u_0(s)S(s)]}{1 - (1-q)s [T(s) - u_0(qs)S(s)]} d_q s \right) d_q t \Big], \end{aligned} \quad (31)$$

$$\begin{aligned} \varphi(x) = & F \exp \left(-\frac{1}{1-q} \int_0^x \frac{\ln \{1 - (1-q)t [T(t) - u_0(qt)S(t)]\}}{t} d_q t \right) + \\ & + u_0(x) \exp \left(-\frac{1}{1-q} \int_0^x \frac{\ln \{1 - (1-q)t [R(t) + u_0(t)S(t)]\}}{t} d_q t \right) \times \\ & \times \left[D + F \int_0^x \frac{S(t)}{1 - (1-q)t [R(t) + u_0(t)S(t)]} \times \right. \\ & \times \exp \left(\frac{1}{1-q} \int_0^t \frac{1}{s} \ln \frac{1 - (1-q)s [R(s) + u_0(s)S(s)]}{1 - (1-q)s [T(s) - u_0(qs)S(s)]} d_q s \right) d_q t \Big], \end{aligned} \quad (32)$$

The constants D and F are related with the initial conditions by:

$$\begin{aligned} D &= \psi(0), \\ F &= -\psi(0) \frac{\varphi_0(0)}{\psi_0(0)} + \varphi(0). \end{aligned} \quad (33)$$

Proof:

In order to prove the formulae (31- 32) and (33) we assume in (12) that $\Lambda'(x)$ is uppertriangular and we apply the q -difference Darboux transform (17) with

$$D(x) = \begin{pmatrix} 1 & 0 \\ -u_0(x) & 1 \end{pmatrix}. \quad (34)$$

The formulae (30) follows now from (17) and from the property of $\Lambda'(x)$ of being uppertriangular. The formulae (31) and (32) are obtained from **Proposition 1** and (33).

QED

As a corollary of **Proposition 2** we obtained the auto-Bäcklund transform for q -difference Riccati equation.

Proposition 3 Let $u_0(x)$ be some special solution of the equation (6). Then the general solution of (6) is given by

$$u^t(x) = (\mathcal{B}_t^+ u_0)(x) = u_0(x) + \frac{t \exp \left(\frac{1}{1-q} \int_0^x \frac{1}{y} \ln \frac{1-(1-q)y[R(y)+u_0(y)S(y)]}{1-(1-q)y[T(y)-u_0(qy)S(y)]} d_q y \right)}{1 + t \int_0^x \frac{S(y)}{1-(1-q)y[R(y)+u_0(y)S(y)]} \exp \left(\frac{1}{1-q} \int_0^y \frac{1}{s} \ln \frac{1-(1-q)s[R(s)+u_0(s)S(s)]}{1-(1-q)s[T(s)-u_0(qs)S(s)]} d_q s \right) d_q y}, \quad (35)$$

where $t \in \mathbb{R}$.

Proof:

The formulae (35) is obtained by substituting (31) and (32) into (7) and putting $t = \frac{F}{D}$.

QED

The q -difference Schrödinger equation

$$(-\partial_q^2 + V(x)) \psi(x) = 0, \quad (36)$$

is a special case of (6) and is obtained by putting $R(x) = T(x) = 0$ and $S(x) = 1$. From the **Proposition 2** we may draw:

Corollary 1 The solution of q -difference Schrödinger equation with the potential

$$V(x) = \partial_q u_0(x) + u_0(x)u_0(qx), \quad (37)$$

is given by

$$\psi(x) = \exp \left(-\frac{1}{1-q} \int_0^x \frac{\ln(1 - (1-q)tu_0(t))}{t} d_q t \right) \times \left[D + F \int_0^x \frac{1}{1 - (1-q)tu_0(t)} \exp \left(\frac{1}{1-q} \int_0^t \frac{1}{s} \ln \frac{1 - (1-q)su_0(s)}{1 + (1-q)su_0(qs)} d_q s \right) d_q t \right]. \quad (38)$$

In the limit of $q \rightarrow 1$ the q -difference equations (5) and (6) tend to their differential counterparts

$$\frac{d}{dx} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} = \begin{pmatrix} R(x) & S(x) \\ V(x) & T(x) \end{pmatrix} \begin{pmatrix} \psi(x) \\ \varphi(x) \end{pmatrix} \quad (39)$$

and

$$V(x) = \frac{d}{dx} u(x) + (R(x) - T(x)) u(x) + S(x) u^2(x), \quad (40)$$

where $u(x)$ is given by (7). The **Propositions 2** and **3** are valid in the limit $q \rightarrow 1$ too. Therefore, we can apply them to the differential case and reproduce in such a way the formulae for general solutions of (39) and (40). They are given by

$$\begin{aligned} \psi(x) &= \exp \left(\int_0^x (R(t) + u_0(t)S(t)) dt \right) \times \\ &\times \left[D + F \int_0^x S(t) \exp \left(\int_0^t [T(s) - R(s) - 2u_0(s)S(s)] ds \right) dt \right], \quad (41) \\ \varphi(x) &= F \exp \left(\int_0^x (T(t) - u_0(t)S(t)) dt \right) + \\ &+ u_0(x) \exp \left(\int_0^x (R(t) + u_0(t)S(t)) dt \right) \times \end{aligned}$$

$$\times \left[D + F \int_0^x S(t) \exp \left(\int_0^t [T(s) - R(s) - 2u_0(s)S(s)] ds \right) dt \right] \quad (42)$$

and by

$$u^t(x) = u_0(x) + \frac{t \exp \left(\int_0^x [T(y) - R(y) - 2u_0(y)S(y)] dy \right)}{1 + t \int_0^x S(y) \exp \left(\int_0^y [T(s) - R(s) - 2u_0(s)S(s)] ds \right) dy}. \quad (43)$$

Here $(\psi_0(x) \ \varphi_0(x))^\top$ and $u_0(x)$ are some special solutions (39) and (40).

In order to describe the properties of the family of solutions $u^t(x) = (\mathcal{B}_t^+ u_0)(x)$, $t \in \mathbb{R}$, given by (35) let us formulate:

Proposition 4

- The transforms (35) form one-parameter group

$$\mathcal{B}_{t_1}^+ \circ \mathcal{B}_{t_2}^+ = \mathcal{B}_{t_1+t_2}^+, \quad (44)$$

which acts transitively on the space of all solutions of the q -difference Riccati equations (6).

- The solutions $u^{t_1}(x)$, $u^{t_2}(x)$, $u^{t_3}(x)$, $u^{t_4}(x)$ satisfy the unharmonic superposition principle

$$\frac{(u^{t_4}(x) - u^{t_3}(x))(u^{t_1}(x) - u^{t_2}(x))}{(u^{t_3}(x) - u^{t_1}(x))(u^{t_2}(x) - u^{t_4}(x))} = \frac{(t_4 - t_3)(t_1 - t_2)}{(t_3 - t_1)(t_2 - t_4)}, \quad (45)$$

for $t_1, t_2, t_3, t_4 \in \mathbb{R}$.

Proof:

- According to **Proposition 3** any solution of (6) is given by $\mathcal{B}_s u_0$ for some $s \in \mathbb{R}$. Since $\mathcal{B}_{t_1}(\mathcal{B}_{t_2} u_0)$ is a solution of (6) we have

$$\mathcal{B}_{t_1}^+ \circ \mathcal{B}_{t_2}^+ u_0 = \mathcal{B}_s^+ u_0. \quad (46)$$

One can find $s \in \mathbb{R}$ by evaluation of both sides of (46) at $x = 0$. Thus, using (33) we obtain

$$t_1 + (t_2 + u_0(0)) = s + u_0(0). \quad (47)$$

- The equality is obtained from (35) by direct calculation.

QED

The right hand side of (45) is invariant with respect to the real fractional transformation

$$t'_i = \frac{at_i + b}{ct_i + d}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R}) \quad (48)$$

of the parameters t_1, t_2, t_3 and t_4 . Hence, the left hand side of (45) is $SL(2, \mathbb{R})$ -invariant too.

3 Some integrable cases

In the **Proposition 3** we constructed the transform which generated the general solution of the q -Riccati equation (6) from a given special one. It was done by the action of one-parameter group of transformations $\{\mathcal{B}_t^+\}$, $t \in \mathbb{R}$. The formula (35) which defines the group $\{\mathcal{B}_t^+\}_{t \in \mathbb{R}}$ action does depend on the potentials $R(x)$, $S(x)$ and $T(x)$. It does not contain the potential $V(x)$. This allows us to define some method of creation of new integrable systems from a system which one knows how to integrate. In order to do this let us introduce some notation.

By $\mathcal{I}$ we will denote the map

$$\mathcal{I}u(x) := -u(x). \quad (49)$$

The operator which acts on the function $u(x)$ on the right side of q -Riccati equation (6) will be denoted by $\mathcal{R}_+$, i.e.

$$\mathcal{R}_+u(x) := \partial_q u(x) - T(x)u(x) + R(x)u(qx) + S(x)u(x)u(qx). \quad (50)$$

By $\mathcal{R}_-$ we will denote the operator

$$\mathcal{R}_-u(x) := -\partial_q u(x) + T(x)u(x) - R(x)u(qx) + S(x)u(x)u(qx). \quad (51)$$

It is clear that

$$\mathcal{R}_+ \circ \mathcal{I} = \mathcal{R}_- \quad \text{and} \quad \mathcal{R}_- \circ \mathcal{I} = \mathcal{R}_+. \quad (52)$$

Hence, if $u(x)$ is the solution of the equation

$$\mathcal{R}_+u(x) = V(x), \quad (53)$$

then $\mathcal{I}u(x)$ does satisfy

$$\mathcal{R}_- \circ \mathcal{I}u(x) = V(x) \quad (54)$$

and vice versa. The one-parameter groups (35) for the equations (53) and (54) will be denoted by $\{\mathcal{B}_t^+\}_{t \in \mathbb{R}}$ and $\{\mathcal{B}_t^-\}_{t \in \mathbb{R}}$ respectively. They are related by

$$\mathcal{B}_t^- = \mathcal{I} \circ \mathcal{B}_t^+ \circ \mathcal{I}. \quad (55)$$

After application of the transform

$$\mathcal{B}_{t_1 \dots t_n}^- := \mathcal{I} \circ \mathcal{B}_{t_1}^+ \circ \mathcal{I} \circ \mathcal{B}_{t_2}^+ \circ \dots \circ \mathcal{I} \circ \mathcal{B}_{t_n}^+ \circ \mathcal{I} \quad (56)$$

to $u_0(x)$ which one assumes to be the solution of (6) with the potential $V_0(x)$ we obtain the solution

$$u(t_1, \dots, t_n, x) := \mathcal{B}_{t_1 \dots t_n}^- u_0(x) \quad (57)$$

of (53) with the some new potential

$$\mathcal{R}_+u(t_1, \dots, t_n, x) = V(t_1, \dots, t_n, x), \quad (58)$$

which is n -parameter deformation of the initial one. The same function satisfies also

$$\mathcal{R}_-u(t_1, \dots, t_n, x) = V(t_2, \dots, t_n, x). \quad (59)$$

In such a way we obtain the family of q -difference Riccati equations (58) and (59) generated by $u_0(x)$ and by the transform (56).

It is worth to mention here the group like property of $\mathcal{B}_{t_1 \dots t_n}$:

$$\begin{aligned} \mathcal{B}_{0 \dots 0}^- &= id, \\ (\mathcal{B}_{t_1 \dots t_n}^-)^{-1} &= \mathcal{B}_{-t_n \dots -t_1}, \\ \mathcal{B}_{t_1 \dots t_n}^- \circ \mathcal{B}_{s_1 \dots s_m}^- &= \mathcal{B}_{t_1 \dots t_{n-1} t_n + s_1 s_2 \dots s_m}^-. \end{aligned} \quad (60)$$

In particular of case $n = 1$, applying $\{\mathcal{B}_t^-\}_{t \in \mathbb{R}}$ to (37) we obtain

$$\mathcal{R}_+ u(t, x) = V(t, x), \quad (61)$$

$$\mathcal{R}_- u(t, x) = \mathcal{R}_- u_0(x), \quad (62)$$

where $u(t, x) = \mathcal{B}_t^- u_0(x)$. Simple calculation gives a solution

$$\begin{aligned} u(t, x) &= u_0(x) - \\ &- \frac{t \exp \left(\frac{1}{1-q} \int_0^x \frac{1}{y} \ln \frac{1+(1-q)y u_0(y)}{1-(1-q)y u_0(qy)} d_q y \right)}{1 + t \int_0^x \frac{1}{1+(1-q)y u_0(y)} \exp \left(\frac{1}{1-q} \int_0^y \frac{1}{s} \ln \frac{1+(1-q)s u_0(s)}{1-(1-q)s u_0(qs)} d_q s \right) d_q y}, \end{aligned} \quad (63)$$

for the potential

$$\begin{aligned} V(t, x) &= V_0(x) - \\ &- 2\partial_q \frac{t \exp \left(\frac{1}{1-q} \int_0^x \frac{1}{y} \ln \frac{1+(1-q)y u_0(y)}{1-(1-q)y u_0(qy)} d_q y \right)}{1 + t \int_0^x \frac{1}{1+(1-q)y u_0(y)} \exp \left(\frac{1}{1-q} \int_0^y \frac{1}{s} \ln \frac{1+(1-q)s u_0(s)}{1-(1-q)s u_0(qs)} d_q s \right) d_q y}. \end{aligned} \quad (64)$$

In the limit of $q \rightarrow 1$ the equations (61) and (62) correspond to the proper differential equations and (63) tends to the solution

$$u(t, x) = u_0(x) - \frac{\partial}{\partial x} \ln \left(1 + t \int_0^x \exp \left(2 \int_0^y u_0(z) dz \right) dy \right) \quad (65)$$

of (3) with potential given by

$$V(t, x) = V_0(x) - 2 \frac{\partial^2}{\partial x^2} \ln \left(1 + t \int_0^x \exp \left(2 \int_0^y u_0(z) dz \right) dy \right). \quad (66)$$

Combining (61) and (62) we find the formula

$$\begin{aligned} u(t, x) &= \frac{1}{2} \left\{ (1-q) x \frac{1}{2} (V(t, x) - \mathcal{R}_- u_0(x)) \right. \\ &\left. \pm \sqrt{\left[(1-q) x \frac{1}{2} (V(t, x) - \mathcal{R}_- u_0(x)) \right]^2 + 2(V(t, x) + \mathcal{R}_- u_0(x))} \right\}, \end{aligned} \quad (67)$$

which expresses the solution $u(t, x)$ by the t -deformed potential $V(t, x)$ and by the initial solution $u_0(x)$.

Example

The function

$$u_0(x) = ax^\alpha, \quad (68)$$

where $\alpha > -1$, $a, \alpha \in \mathbb{R}$, is the solution of (37) with the potential

$$V_0(x) = a \frac{1-q^\alpha}{1-q} x^{\alpha-1} + a^2 q^\alpha x^{2\alpha}. \quad (69)$$

Let

$$\exp_R(x) = \sum_{n=0}^{\infty} \frac{1}{R(q) \cdots R(q^n)} x^n, \quad (70)$$

be generalized exponential function (see [3]) with

$$R(x) = \frac{1 - x^{\alpha+1}}{(1 - q) x^{\alpha+1}}. \quad (71)$$

Using the formulae (63) and (64) we find that

$$u(t, x) = ax^{\alpha} - \frac{t \frac{\exp_R(ax^{\alpha+1})}{\exp_R(-aq^{\alpha}x^{\alpha+1})}}{1 + t \int_0^x \frac{\exp_R(aq^{\alpha+1}y^{\alpha+1})}{\exp_R(-aq^{\alpha}y^{\alpha+1})} d_q y}, \quad (72)$$

is a solution of (37) with potential $V(t, x)$ given by

$$\begin{aligned} V(t, x) = & a \frac{1 - q^{\alpha}}{1 - q} x^{\alpha-1} + a^2 q^{\alpha} x^{2\alpha} - \\ & - 2t \frac{\frac{\exp_R(aq^{\alpha+1}x^{\alpha+1})}{\exp_R(-aq^{\alpha}x^{\alpha+1})}}{\left(1 + t \int_0^x \frac{\exp_R(aq^{\alpha+1}y^{\alpha+1})}{\exp_R(-aq^{\alpha}y^{\alpha+1})} d_q y\right) \left(1 + qt \int_0^x \frac{\exp_R(aq^{2(\alpha+1)}y^{\alpha+1})}{\exp_R(-aq^{2\alpha+1}y^{\alpha+1})} d_q y\right)} \times \\ & \times \left\{ (1 + q^{\alpha}) a q^{\alpha+1} x^{\alpha} \left(1 + t \int_0^x \frac{\exp_R(aq^{\alpha+1}y^{\alpha+1})}{\exp_R(-aq^{\alpha}y^{\alpha+1})} d_q y\right) - t \frac{\exp_R(aq^{\alpha+1}x^{\alpha+1})}{\exp_R(-aq^{2\alpha+1}x^{\alpha+1})} \right\}. \quad (73) \end{aligned}$$

If, for example, $\alpha = 0$, $t = a$ and $q \rightarrow 1$ then the potential (73) reduces to Rosen–Morse potential

$$V(a, x) = a^2 \frac{\exp(-2ax) - 6 + \exp(2ax)}{(\exp(-ax) + \exp(ax))^2}. \quad (74)$$

Acknowledgements

We thank dr Z Hasiewicz for helpful comments.

References

- [1] V.B. Matveev, M.A. Salle "Darboux Transformations and Solitons", Springer-Verlag, Berlin Heidelberg, 1991;
- [2] S.P. Novikov, I.A. Dynnikov "Discrete spectral symmetries of low-dimensional differential operators and difference operators on regular lattices and two-dimensional manifolds", Russian Math. Surveys 52 (1997), no.5, 1057–1116;
- [3] A. Odziejewicz "Quantum Algebras and q -Special Functions Related to Coherent States Maps of the Disc", Commun. Math. Phys. 192, 183-215 (1998);
- [4] I.K. Sujetin "Classical orthogonal polynomials", "Nauka", (in Russian), Moscow 1979;
- [5] B. Mielnik, L.M. Nieto, O. Rosas-Ortiz "The finite difference algorithm for higher order supersymmetry", Physics Letters A 269 (2000) 70-78;

- [6] J.F. Carinena, A. Ramos "Shape-invariant potentials depending on n parameters transformed by translation", J. Phys. A: Math. Gen. 33 (2000) 3467-3481;
- [7] O.L. De Lange, R.E. Raab "Operator method in quantum mechanics", Oxford Science Publication, Calderon Press, Oxford, 1991;
- [8] W. Miller Jr. "Lie theory and Special Functions", Academic Press, New York and London, 1968;
- [9] I.M. Ryžyk, I.C. Gradsztejn "The tables of integrals, sums, series and products", GITTL, (in Russian), Moscow 1951;